

DISEASE	USUAL INCUBATION PERIOD	PERIOD OF INFECTIOUSNESS	EXCLUSION PERIOD
CHICKENPOX AND SHINGLES	13-17 days	1-2 days before rash appears until 5 days after rash (7 days shingles)	5- 10 days after last spot appears, and until all the spots have crusted over.
CONJUNCTIVITIS	3-5 days		48 hours after first drops have been given
COUGHS AND COLDS	2-3 days	Just before and for 3 days after onset of symptoms	Until child fully recovered
COXSACKIE(HAND, FOOT AND MOUTH)	3-6 days	Whilst virus is present in faeces - For about 4 weeks after the child has recovered	Until the spots and sores have gone and child feels well
COVID	2-14 days	2 weeks	They no longer have a high temperature and feel well enough to attend. Follow the national guidance if they've tested positive for COVID-19
FIFTH DISEASE (SLAPPED CHEEK)	Variable 4-20 days	Until child feels well	Until child fully recovered
GASTRO-ENTERITIS SICKNESS & DIARRHOEA	Varies-a few hours to a few days	Until sickness/diarrhoea stops	Until 48 hours after sickness/diarrhoea stops
GLANDULAR FEVER	4-6 weeks	While virus is present in saliva	Until clinically recovered
HEAD LICE	Eggs hatch and become mature in 1 - 2 weeks	Whilst lice or viable eggs are present	Until commencement of treatment
HEPITITIS A INFECTIOUS	14-42 days	14 days before jaundice to 7 days after	7 days from jaundice starting and child feels well
HEPATITIS B,C,HIV			None
HERPES SIMPLEX	2-12 days		No need to exclude but avoid contact at all times with sores
INFLUENZA (Flu)	2-3 days	Just before and for 3 days after onset of symptoms	Until child is fully recovered
MEASLES	7-14 days	1 day before first symptoms until 4 days after onset of rash	4 days after onset of rash

MENINGOCOCCAL MENINGITIS/SEPTICAEMIA			Until recovered, at least 48 hours after treatment has started
MENINGITIS due to other bacteria			Until recovered, at least 48 hours after treatment has started
MENINGITIS VIRAL			Until recovered, at least 48 hours after treatment has started
MRSA			Until recovered, at least 48 hours after treatment has started
MUMPS	18 days	7 days before swelling until 9 days after	5 days after onset of swelling
RINGWORM			48 hours after treatment has started
RUBELLA (GERMAN MEASLES)	16-18 days	7 days before rash until 4 days after	Until 6 days after rash appears. Pregnant staff or parent contacts should seek prompt advice from their GP or midwife.
SCABIES			48 hours after treatment has started
STREPTOCOCCAL INFECTION(SCARLET FEVER)	1-3 DAYS	Day sore throat starts until 24 hours after starting antibiotics	Until clinically recovered or antibiotic treatment
THREADWORMS			48 hours after treatment has started (treatment also recommended for child's household)
TONSILLITIS			48 hours after treatment has started
TUBERCULOSIS (TB)			48 hours after treatment has started
STREPTOCOCCAL INFECTION(IMPETIGO)	1-3 days	Until the lesions are healed or until 48 hours after the start of effective treatment	Until sores are healed or until 48 hours after antibiotic treatment
WHOOPING COUGH	7-10 days	2-4 days before cough until 21 days afterwards.	2 days from commencing the antibiotic treatment or 21 days if no antibiotic treatment given

CORONAVIRUS	Up to 10 days advisory	Daily updates for advisories noting the category of the countries you visited	Between 1 & 10 days of diagnosis with self-isolation and given the all clear.
-------------	------------------------	---	---

**** If your child has been prescribed antibiotics for ANY reason they must stay away from nursery for at least 48 hours.**